

# Energy dependence of hadronic observables in central Pb+Pb reactions at the CERN SPS

Christoph Blume<sup>1</sup> for the NA49 Collaboration<sup>a</sup>

<sup>1</sup> Institut für Kernphysik der J.W. Goethe Universität,  
60486 Frankfurt am Main, Germany

## Abstract.

*Keywords:* Spectra, Fluctuations, HBT


*PACS:* 25.75Nq

## 1. Introduction

In the recent years the NA49 experiment has collected data on Pb+Pb collisions at beam energies between 20 to 158 AGeV with the objective to cover the critical region of energy densities where the expected phase transition to a deconfined phase might occur in the early stage of the reactions. In this contribution the energy dependence of various hadronic observables is presented. These include  $m_t$ - and rapidity-distributions, particle ratios and particle ratio fluctuations, as well as HBT radii. NA49 is a fixed target experiment at the CERN SPS and consists of a large acceptance magnetic spectrometer equipped with four TPCs as tracking devices and a forward calorimeter for centrality selection. Details on the experimental setup can be found in [ 1].

## 2. Particle spectra

Characterizing the energy dependence of the shape of  $m_t$ -spectra generally requires a comparison to a model. A simple exponential fit to the data can be misleading, since most particle spectra exhibit a clear curvature. Pion  $m_t$  spectra are concave, mainly due to resonance contributions, while proton spectra measured in heavy ion reactions are convex, caused by the effect of radial flow. To a certain extent, kaons are an exception that can relatively well be approximated by an exponential. An analysis of the slope parameter for kaons, extracted by an exponential fit, revealed a clear change of its energy dependence around beam energies of 20-30 AGeV [ 2]. Figure 1 summarizes the energy dependence of  $\langle m_t \rangle - m_0$ . This quantity has the advantage of providing a model independent characterization of the transverse mass




**Fig. 1.** The  $\sqrt{s}$ -dependence of  $\langle m_t \rangle - m_0$  for pions, kaons, and protons at mid-rapidity for 5-10% most central Au+Au/Pb+Pb reactions.

distributions and can therefore easily be employed for pions and protons as well. As can be seen, the change in the evolution of the  $m_t$ -spectra with energy is clearly also present for pions, similar to the kaons, and in a less pronounced fashion for the protons as well.

The large acceptance of the NA49 spectrometer allows to measure particle spectra over a wide range of the longitudinal phase space. Figure 2 shows a compilation of the rapidity distributions of  $\pi^-$ ,  $K^+$ ,  $K^-$ ,  $\phi$ , and  $\Lambda$  [ 2]. Generally, a clear increase of the widths with beam energy can be observed, where the width of the  $\pi^-$  distribution is approximately equal to the  $K^+$  and both are wider than the  $K^-$  distribution. The shape of the distribution for pions and kaons is well described by a Gaussian. The  $\Lambda$ -distributions, however, exhibit a strong variation of the shape: While at 30 AGeV they are still Gaussian-shaped, a clear plateau develops with increasing beam energy.

### 3. Particle multiplicities

The rapidity distributions, discussed above, allow to determine the total yields of the different particle species with only small extrapolations. Fig.3 shows the ratio of the resulting  $4\pi$ -yields of K and  $\pi$  [ 2, 3]. While this ratio for negatively charged particles rises more or less continuously (left hand side of Fig. 3) – except a small indication for a kink at 30 AGeV – a very distinct maximum is observed in the positively charged case (right hand side of Fig. 3). The lines included in the figures are predictions of an extended hadron gas model [ 4] and the transport codes RQMD [ 5] and UrQMD [ 6]. Even though the hadron gas model and RQMD predict a maximum of the  $\langle K^+ \rangle / \langle \pi^+ \rangle$ -ratio in the SPS energy range, none of the models can fully describe the sharp feature of its energy dependence. A feature, which is also



**Fig. 2.** The rapidity spectra of hadrons produced in central (7% at 20-80 AGeV, 5% ( $\pi^-$ ,  $K^+$ ,  $K^-$ ) and 10% ( $\phi$ ,  $\Lambda$ ) at 158 AGeV. The closed symbols indicate measured points, open points are reflected with respect to mid-rapidity. The solid lines indicate parametrizations of the data used for the extrapolation of the yield to full phase space.


not present in p+p collisions. It is also noteworthy that the hadron gas model does not fit the  $\langle K^- \rangle / \langle \pi^- \rangle$ -ratio at energies above 40 AGeV either. On the other hand, a strong non-monotonic energy dependence of the total strangeness to pion ratio was predicted by the Statistical Model of The Early Stage [7], assuming a phase transition from confined matter to a quark-gluon plasma in the SPS energy range.

#### 4. Particle ratio fluctuations

NA49 has performed an event-by-event measurement of the particle ratios [8]. The dynamical fluctuations  $\sigma_{dyn}$  of this ratios have been extracted by subtracting the r.m.s. width  $\sigma_{mix}$  of the mixed event distributions from the r.m.s. width  $\sigma_{data}$  of the real event distributions:

$$\sigma_{dyn} = \text{sign}(\sigma_{data}^2 - \sigma_{mixed}^2) \sqrt{|\sigma_{data}^2 - \sigma_{mixed}^2|} \quad (1)$$

The mixed event particle ratios contain by construction the effects of finite number fluctuations as well as effects of the detector resolution. As shown in the left panel



**Fig. 3.** The energy dependence of the  $\langle K \rangle / \langle \pi \rangle$  ratios together with various model predictions (see text).

of Fig. 4, the  $K/\pi$  fluctuations are positive and decrease with beam energy. The  $p/\pi$  fluctuations, on the other hand, are negative – indicating a correlation present in the real data – and increase with beam energies. While the trend of the  $K/\pi$  fluctuations is not reproduced by UrQMD [9], it provides a good description of the energy dependence of the  $p/\pi$  fluctuations. This might indicate that the negative value of the fluctuations in this ratio is due to resonance decays.

## 5. Bose-Einstein correlations


Figure 5 summarizes the HBT-radii extracted in the LCMS, as measured by the NA49 experiment [10]. As expected in the presence of longitudinal and transverse expansion, a significant reduction of the radii with increasing  $k_t$  is observed at all beam energies. Assuming a boost-invariant scenario, the  $k_t$ -dependence of  $R_{long}$  should reflect the life time of the source [11]:

$$R_{long} = \tau_f \left( \frac{T_f}{m_t} \right)^{1/2} ; \quad m_t = (m_\pi^2 + k_t^2)^{1/2} \quad (2)$$

The fits of this function, assuming a freeze-out temperature  $T_f = 120$  MeV, are shown in the upper part of Fig. 5. Only a weak increase of the extracted life time with beam energy is observed. Another important feature of this data is the fact that  $R_{out} > R_{side}$  at all beam energies (lower part of Fig. 5). The difference of these two parameter is connected to the emission duration [12]:

$$\Delta\tau^2 = \frac{1}{\beta_t^2} (R_{out}^2 - R_{side}^2) ; \quad \beta_t \approx \frac{k_t}{m_t} \quad (3)$$

The data would indicate an emission duration of 3-4 fm/c.



**Fig. 4.** Energy dependence of the event-by-event fluctuation signal of the  $(K^+ + K^-)/(\pi^+ + \pi^-)$  ratio (left hand side) and the  $(p + \bar{p})/(\pi^+ + \pi^-)$  ratio (right hand side). The systematic errors of the measurements are shown as gray bands.

## 6. Conclusions

The recent study of the excitation functions of hadronic observables in the SPS energy range has revealed a number of interesting and unexpected results. This includes a step-like energy dependence of the  $\langle m_t \rangle - m_0$  of pions and kaons and a sharp maximum in the strangeness to pion ratio. The dynamical  $K/\pi$  ratio fluctuations are positive and decrease with beam energy in the range between 20 - 158 AGeV, while the  $p/\pi$  ratio fluctuations are negative and increase. The HBT-radii, however, do not exhibit a significant energy dependence in this energy range.


## Acknowledgements

This work was supported by the US Department of Energy Grant DE-FG03-97ER41020/A000, the Bundesministerium für Bildung und Forschung, Germany, the Polish State Committee for Scientific Research (2 P03B 130 23, SPB/CERN/P-03/Dz 446/2002-2004, 2 P03B 04123), the Hungarian Scientific Research Foundation (T032648, T032293, T043514), the Hungarian National Science Foundation, OTKA, (F034707), the Polish-German Foundation, and the Korea Research Foundation Grant (KRF-2003-070-C00015).

## Notes

a. The NA49 collaboration:

C. Alt<sup>9</sup>, T. Anticic<sup>21</sup>, B. Baatar<sup>8</sup>, D. Barna<sup>4</sup>, J. Bartke<sup>6</sup>, L. Betev<sup>9,10</sup>, H. Bialkowska<sup>19</sup>,



**Fig. 5.** The HBT-radii as a function of  $k_t$  at mid-rapidity for central Pb+Pb collisions at 20 to 158 AGeV

A. Billmeier<sup>9</sup>, C. Blume<sup>9</sup>, B. Boimska<sup>19</sup>, M. Botje<sup>1</sup>, J. Bracinik<sup>3</sup>, R. Bramm<sup>9</sup>, R. Brun<sup>10</sup>, P. Buncić<sup>9,10</sup>, V. Cerny<sup>3</sup>, P. Christakoglou<sup>2</sup>, O. Chvala<sup>15</sup>, J.G. Cramer<sup>17</sup>, P. Csató<sup>4</sup>, N. Dardenov<sup>18</sup>, A. Dimitrov<sup>18</sup>, P. Dinkelaker<sup>9</sup>, V. Eckardt<sup>14</sup>, G. Farantatos<sup>2</sup>, D. Flierl<sup>9</sup>, Z. Fodor<sup>4</sup>, P. Foka<sup>7</sup>, P. Freund<sup>14</sup>, V. Friese<sup>7</sup>, J. Gál<sup>4</sup>, M. Gaździcki<sup>9</sup>, G. Georgopoulos<sup>2</sup>, E. Gładysz<sup>6</sup>, K. Grebieszko<sup>20</sup>, S. Hegyi<sup>4</sup>, C. Höhne<sup>13</sup>, K. Kadija<sup>21</sup>, A. Karev<sup>14</sup>, M. Kliemant<sup>9</sup>, S. Kniese<sup>9</sup>, V.I. Kolesnikov<sup>8</sup>, T. Kollegger<sup>9</sup>, E. Kornas<sup>6</sup>, R. Korus<sup>12</sup>, M. Kowalski<sup>6</sup>, I. Kraus<sup>7</sup>, M. Kreps<sup>3</sup>, M. van Leeuwen<sup>1</sup>, P. Lévai<sup>4</sup>, L. Litov<sup>18</sup>, B. Lungwitz<sup>9</sup>, M. Makariev<sup>18</sup>, A.I. Malakhov<sup>8</sup>, C. Markert<sup>7</sup>, M. Mateev<sup>18</sup>, B.W. Mayes<sup>11</sup>, G.L. Melkumov<sup>8</sup>, C. Meurer<sup>9</sup>, A. Mischke<sup>7</sup>, M. Mitrovski<sup>9</sup>, J. Molnár<sup>4</sup>, S. Mrówczyński<sup>12</sup>, G. Pál<sup>4</sup>, A.D. Panagiotou<sup>2</sup>, D. Panayotov<sup>18</sup>, A. Petridis<sup>2</sup>, M. Pikna<sup>3</sup>, L. Pinsky<sup>11</sup>, F. Pühlhofer<sup>13</sup>, J.G. Reid<sup>17</sup>, R. Renfordt<sup>9</sup>, A. Richard<sup>9</sup>, C. Roland<sup>5</sup>, G. Roland<sup>5</sup>, M. Rybczyński<sup>12</sup>, A. Rybicki<sup>6,10</sup>, A. Sandoval<sup>7</sup>, H. Sann<sup>7</sup>, N. Schmitz<sup>14</sup>, P. Seyboth<sup>14</sup>, F. Siklér<sup>4</sup>, B. Sitar<sup>3</sup>, E. Skrzypczak<sup>20</sup>, G. Stefanek<sup>12</sup>, R. Stock<sup>9</sup>, H. Ströbele<sup>9</sup>, T. Susa<sup>21</sup>, I. Szentpétery<sup>4</sup>, J. Sziklai<sup>4</sup>, T.A. Trainor<sup>17</sup>, V. Trubnikov<sup>20</sup>, D. Varga<sup>4</sup>, M. Vassiliou<sup>2</sup>, G.I. Veres<sup>4,5</sup>, G. Vesztergombi<sup>4</sup>, D. Vranić<sup>7</sup>, A. Wetzler<sup>9</sup>, Z. Włodarczyk<sup>12</sup>, I.K. Yoo<sup>16</sup>, J. Zaranek<sup>9</sup>, J. Zimányi<sup>4</sup>

<sup>1</sup>NIKHEF, Amsterdam, Netherlands.

<sup>2</sup>Department of Physics, University of Athens, Athens, Greece.

<sup>3</sup>Comenius University, Bratislava, Slovakia.

<sup>4</sup>KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary.

<sup>5</sup>MIT, Cambridge, USA.

<sup>6</sup>Institute of Nuclear Physics, Cracow, Poland.

<sup>7</sup>Gesellschaft für Schwerionenforschung (GSI), Darmstadt, Germany.

<sup>8</sup>Joint Institute for Nuclear Research, Dubna, Russia.

<sup>9</sup>Fachbereich Physik der Universität, Frankfurt, Germany.

- <sup>10</sup>CERN, Geneva, Switzerland.
- <sup>11</sup>University of Houston, Houston, TX, USA.
- <sup>12</sup>Institute of Physics Świetokrzyska Academy, Kielce, Poland.
- <sup>13</sup>Fachbereich Physik der Universität, Marburg, Germany.
- <sup>14</sup>Max-Planck-Institut für Physik, Munich, Germany.
- <sup>15</sup>Institute of Particle and Nuclear Physics, Charles University, Prague, Czech Republic.
- <sup>16</sup>Department of Physics, Pusan National University, Pusan, Republic of Korea.
- <sup>17</sup>Nuclear Physics Laboratory, University of Washington, Seattle, WA, USA.
- <sup>18</sup>Atomic Physics Department, Sofia University St. Kliment Ohridski, Sofia, Bulgaria.
- <sup>19</sup>Institute for Nuclear Studies, Warsaw, Poland.
- <sup>20</sup>Institute for Experimental Physics, University of Warsaw, Warsaw, Poland.
- <sup>21</sup>Rudjer Boskovic Institute, Zagreb, Croatia.

## References

1. S. V. Afanasiev et al., Nucl. Instrum. Meth. A430 (1999) 210-244.
2. M. Gaździcki (for the NA49 collaboration), arXiv: nucl-ex/0403023.
3. S. V. Afanasiev et al., Phys. Rev. C66 (2002) 157-166.
4. P. Braun-Munzinger, K. Redlich, and H. Oeschler, Nucl. Phys. A697 (2002) 902-912.
5. H. Sorge, H. Stöcker, and W. Greiner, Nucl. Phys. A498 (1989) 567.
6. S. A. Bass et al., Prog. Part. Nucl. Phys. 41 (1998) 255-369.
7. M. Gazdzicky and M. I. Gorenstein, Acta Phys. Polon. B30 (1999) 2705-2735.
8. C. Roland (for the NA49 collaboration), arXiv: nucl-ex/0403035.
9. M. Bleicher et al., J. Phys. G25 (1999) 1859.
10. S. Kniege (for the NA49 collaboration), arXiv: nucl-ex/0403034.
11. M. Makhlin and Y.M. Synukov, Z. Phys. C39 (1988) 2694.
12. U. Heinz et al., Phys. Lett. B382 (1996) 181.